
LETTERS
TO THE EDITOR

Dedicated to V. F. Mironov on His 60th Anniversary

Synthesis, Acid-Base and Complexing Properties of *N,N,N',N'*-Tetrakis(*O*-butylhydroxyphosphorylmethyl)- 1,2-diaminoethane

I. D. Shurygin^{a*}, A. R. Garifzyanov^a, R. A. Cherkasov^a, K. A. Ivshin^{a,b}, and O. N. Kataeva^{a,b}

^a Kazan Federal University, ul. Kremlevskaya 18, Kazan, Tatarstan, 420008 Russia

*e-mail: idshurygin@gmail.com

^b Arbuzov Institute of Organic and Physical Chemistry, Kazan Scientific Center,
Russian Academy of Sciences, Kazan, Tatarstan, Russia

Received May 11, 2017

Abstract—A method of synthesis of organophosphorus analog of ethylenediaminetetraacetic acid (a common complexon), *N,N,N',N'*-tetrakis(*O*-butylhydroxyphosphorylmethyl)-1,2-diaminoethane, has been elaborated. The product structure has been elucidated by means of NMR spectroscopy and X-ray diffraction analysis. Constants of the acid dissociation and the stability of the 1 : 1 complexes with Co(II), Ni(II), and Cu(II) ions have been determined.

Keywords: organophosphorus complexon, acid-base properties, complex stability constant

DOI: 10.1134/S1070363217090274

We have earlier synthesized an organophosphorus analog of nitrilotriacetic acid (a common complexon), potassium salt of aminotris(*O*-butylmethylphosphonic) acid [1], and studied its acid-base properties as well as the complex formation with certain doubly charged transition metals cations [2, 3]. We have revealed that this triphosphorylated amine is significantly different from nitrilotriacetic acid in the complex formation, yielding weaker complexes.

In this study, we suggested a method for the synthesis of an organophosphorus analog of another complexon widely used in analytical and coordination chemistry (ethylenediaminetetraacetic acid EDTA): *N,N,N',N'*-tetrakis(*O*-butylhydroxyphosphorylmethyl)-1,2-diaminoethane **4**. On top of that, acid-base and complex-forming properties of the latter were investigated. The interaction of 1,3,6,8-tetraazatricyclodecane (synthesized via α -condensation of formaldehyde with ethylenediamine [4]) afforded earlier unknown *N,N,N',N'*-tetrakis(butoxymethyl)ethane-1,2-diamine **1**. The reaction of the latter with 4 eq. of di-butyl phosphonate led to the formation of the tetraphosphorylated diamine **2** (with almost quanti-

tative yield, ³¹P NMR monitoring) that was transformed without purification into readily crystallizing tetrapotassium salt **3** via hydrolysis with potassium hydroxide. The addition of nitric acid to a solution of salt **3** gave acid **4** (Scheme 1).

According to X-ray diffraction data, acid **4** existed in the crystal in the form of zwitterion salt: both nitrogen atoms were protonated and two hydroxy groups at the phosphorus atom were deprotonated (Fig. 1). The negative charge was localized at the O³ and O^{3'} oxygen atoms, the bond lengths in the deprotonated phosphorus-containing moiety being P¹–O³ 1.516(2), P¹–O¹ 1.466(2), and P¹–O² 1.575(2) Å and those in the other phosphoryl fragment being P²–O⁴ 1.480(2), P²–O⁵ 1.510(2), and P²–O⁶ 1.585(2) Å. The molecule was centrosymmetrical, being in a special position of the *P*-1 space group. The N–C bonds were in the *gauche* positions with respect to the phosphoryl groups. The orientation of butyl substituents was different: that at the P¹ atom was *gauche*-oriented with respect to the phosphoryl bond, whereas the alkyl group at the P² atom was *trans*-oriented.